

**GN-232**

103612

V Semester B.A./B.Sc. Examination, December - 2019  
(CBCS) (F+R) 2016-17 and Onwards)

**MATHEMATICS - V**

Time : 3 Hours

Max. Marks : 70

**Instruction :** Answer **all** questions.**PART - A**Answer **any five** questions.

1. (a) Give an example of
  - (i) a ring with zero divisor
  - (ii) a non-commutative ring with unity
- (b) In a ring  $(R, +, \cdot)$  prove that  $a \cdot (b - c) = a \cdot b - a \cdot c \quad \forall a, b, c \in R$ .
- (c) Define principal and maximal ideals of a ring  $R$ .
- (d) Find the maximum directional derivative of  $\phi = x^3 y^2 z$  at the point  $(1, -2, 3)$ .
- (e) If  $\vec{f} = 3x^2 \hat{i} + 5xy^2 \hat{j} + xyz^3 \hat{k}$  then, find  $\text{div } \vec{f}$  at  $(1, 2, 3)$ .
- (f) Evaluate :  $\Delta^4 (1 - ax)(1 - bx)(1 - cx)(1 - dx)$ .
- (g) Write Lagrange's Interpolation formula for unequal intervals.
- (h) Using Trapezoidal rule, evaluate  $\int_0^6 f(x) dx$  given :

|        |       |       |       |       |       |       |       |
|--------|-------|-------|-------|-------|-------|-------|-------|
| $x$    | 0     | 1     | 2     | 3     | 4     | 5     | 6     |
| $f(x)$ | 0.146 | 0.161 | 0.176 | 0.190 | 0.204 | 0.217 | 0.230 |

**PART - B**Answer **two** full questions.**2x10=20**

2. (a) Prove that the set  $R = \{0, 1, 2, 3, 4, 5\}$  is a commutative ring w.r.t.  $\oplus_6$  and  $\otimes_6$  as two compositions.
- (b) Prove that a ring  $R$  is without zero divisors if and only if the cancellation laws hold in it.

**OR****P.T.O.**





3. (a) Prove that the necessary and sufficient conditions for a non-empty subset  $S$  to be a subring of  $R$ , are :
- (i)  $S + (-S) = S$  (ii)  $SS \subseteq S$
- (b) Define the right and left ideals of a ring  $R$ . Show that the subset

$$S = \left\{ \begin{pmatrix} a & 0 \\ b & 0 \end{pmatrix} \mid a, b \in \mathbb{Z} \right\} \text{ of } M_2(\mathbb{Z}) \text{ is a left ideal but not a right ideal of } M_2(\mathbb{Z}).$$

4. (a) (i) If 'a' is an element of a commutative ring  $R$ , then prove that  $aR = \{ar \mid r \in R\}$  is an ideal of 'R'.
- (ii) If  $I$  is an ideal of a ring 'R' with unity and  $1 \in I$  then prove that  $I = R$ .
- (b) Find all the principal ideals of the ring  $R = \{0, 1, 2, 3, 4, 5\}$  w.r.t.  $\oplus_6$  and  $\otimes_6$  as two compositions.

OR

5. (a) If  $f: R \rightarrow R'$  is a homomorphism of a ring  $R$  into  $R'$  then prove that
- (i)  $f(0) = 0'$  where  $0$  and  $0'$  are the zero elements of  $R$  and  $R'$  respectively.
- (ii)  $f(-a) = -f(a) \forall a \in R$ .
- (b) State and prove fundamental theorem of homomorphism of rings.

PART - C

Answer **two** full questions.

6. (a) Find the constants  $a$  and  $b$  so that the surfaces  $x^2 + ayz = 3x$  and  $bx^2y + z^3 = (b-8)y$  intersect orthogonally at the point  $(1, 1, -2)$ .
- (b) If  $\phi$  is a scalar point function and  $\vec{f}$  is a vector point function then

2x10=20

$$\text{prove that } \text{div}(\phi \vec{f}) = \phi(\text{div } \vec{f}) + (\text{grad } \phi) \cdot \vec{f}$$

OR

7. (a) If  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  then prove that  $\nabla^2(r^3 \vec{r}) = 18r \vec{r}$  where  $r = |\vec{r}|$

(b) If  $\vec{F} = \nabla(x^3 + y^3 + z^3 - 3xyz)$  then find  $\nabla \cdot \vec{F}$  and  $\nabla \times \vec{F}$

8. (a) Show that  $\text{div}(\vec{a} \times (\vec{r} \times \vec{a})) = 2|\vec{a}|^2$  where  $\vec{a}$  is a constant vector.

- (b) Show that  $\vec{F} = (x^2 - yz)\hat{i} + (y^2 - xz)\hat{j} + (z^2 - xy)\hat{k}$  is irrotational. Also find a scalar function  $\phi$  such that  $\vec{F} = \nabla\phi$

OR







9. (a) (i) If  $\vec{F} = 3xy\hat{i} + 20yz^2\hat{j} - 15xz\hat{k}$  and  $\phi = xyz$ , then find  $\text{curl}(\phi\vec{F})$ .
- (ii) Show that  $\vec{F} = 2x^2z\hat{i} - 10xyz\hat{j} + 3xz^2\hat{k}$  is solenoidal.
- (b) Find  $\text{curl}(\text{curl}\vec{F})$  if  $\vec{F} = x^2y\hat{i} - 2xz\hat{j} + 2yz\hat{k}$ .

**PART - D**Answer **two** full questions.**2x10=20**

10. (a) Use the method of separation of symbols to prove that

$$u_0 - u_1 + u_2 - u_3 + \dots + \infty = \frac{1}{2}u_0 - \frac{1}{4}\Delta u_0 + \frac{1}{8}\Delta^2 u_0 - \frac{1}{16}\Delta^3 u_0 + \dots$$

- (b) Obtain a function whose first difference is
- $x^3 + 3x^2 + 5x + 12$
- .

**OR**

11. (a) Find the number of students from the following data who secured marks not more than 45.

| Marks              | 30 - 40 | 40 - 50 | 50 - 60 | 60 - 70 | 70 - 80 |
|--------------------|---------|---------|---------|---------|---------|
| Number of Students | 35      | 48      | 70      | 40      | 22      |

- (b) Find a polynomial of lowest degree which assumes the values 10, 4, 40, 424, 620 at
- $x = -2, 1, 3, 7$
- and 8 respectively, using Newton's divided difference formula.

12. (a) By employing Newton-Gregory backward difference formula, find
- $f(9.7)$
- from the following data.

|        |    |     |    |     |    |
|--------|----|-----|----|-----|----|
| $x$    | 8  | 8.5 | 9  | 9.5 | 10 |
| $f(x)$ | 50 | 57  | 64 | 71  | 75 |

- (b) Using Simpson's
- $\frac{1}{3}$
- <sup>rd</sup>
- rule, Evaluate
- $\int_0^1 \frac{1}{1+x^2} dx$
- dividing the interval (0, 1) into 8 equal parts.

**OR**

13. (a) Applying Lagrange's formula find
- $f(5)$
- , given that
- $f(1)=2, f(2)=4, f(3)=8$
- and
- $f(7)=128$
- .

- (b) Using Simpson's
- $\frac{3}{8}$
- <sup>th</sup>
- rule, Evaluate
- $\int_4^{5.2} \log_e x dx$
- taking
- $h=0.2$
- .

**- o o o -**